

## **X-ray astronomy: towards the next 50 years!**

*Milan, October 1-5, 2012*

*editors: G. Trinchieri, R. Della Ceca, T. Maccacaro, L. Maraschi, G. Pareschi*

### **TABLE OF CONTENTS**

<i>Index</i>	457
<i>Foreword</i>	464
<b>Session I: history and legacy of X-ray astronomy</b>	
R. Giacconi <i>Considerations on X-ray astronomy</i>	472
Y. Tanaka <i>My early days in X-ray astronomy</i>	485
J. Trümper <i>The history of X-ray astronomy in Germany</i>	493
K. Pounds, M. Watson <i>UK contributions to the development of X-ray astronomy as a major international discipline</i>	501
J. Bleeker, F. Verbunt <i>From zone plate to microcalorimeter</i>	512
L. Piro <i>BeppoSAX: history and legacy</i>	522
<b>Session II: stars, neutron stars, pulsars and supernova remnants</b>	
J.H.M.M. Schmitt <i>50 (38) years of stellar X-ray astronomy</i>	532
P. Zhao, J. E. Grindley, J. Hong, M. Servillat, M. van den Berg <i>Cataclysmic variables discovered in the Chandra Multi-wavelength Plane Surveys</i>	540
S. Mereghetti, N. La Palombara, P. Esposito, A. Tiengo <i>A new class of X-ray sources: hot subdwarfs with compact companions</i>	543

A. Odendaal, P. J. Meintjes, P. A. Charles, A. F. Rajoelimanana <i>X-ray periodicity in the supersoft X-ray sources CAL 83 and SMC 13</i>	545
K. Makishima <i>From Sco X-1 to magnetars</i>	547
N. Rea <i>Neutron stars with huge magnetic storms</i>	554
Y.E. Nakagawa, K. Makishima, T. Enoto, T. Sakamoto, T. Mihara, M. Sugizaki, K. Yamaoka, K. Hurley, A. Yoshida, P. Gandhi, M. Tashiro, M. Morii <i>Wide-band spectra of magnetar burst and persistent emission</i>	560
G.L. Israel <i>Monitoring magnetar outbursts</i>	564
M. Marelli, A. De Luca, P. Caraveo <i>The X-ray properties of the Fermi/LAT pulsars</i>	568
D. Salvetti, A. De Luca, P. A. Caraveo, M. Marelli <i>Completing the census of Fermi pulsars with X-ray observations</i>	572
H. Yamaguchi <i>Spectroscopic studies of iron emissions from supernova remnants with Suzaku</i>	574
Y. Hanabata, M. Sawada, H. Katagiri, A. Bamba, Y. Fukazawa <i>Detailed X-ray study of the supernova remnant W51C with Suzaku</i>	580
M. C. Weisskopf <i>Monitoring the Crab Nebula with Chandra</i>	582
D. Dewey <i>SN 1987A at high resolution</i>	586
K.P. Levenfish, A. M. Bykov, M. Durant, O.Y. Kargaltsev, Y. A. Kropotina, G. G. Pavlov, A. M. Krassilchtchikov, Y. A. Uvarov <i>Finest persistent structures in the Vela PWN</i>	588
Y. Terada, A. Harayama, T. Hayashi, T. Dogtani, M. Ishida, A. Bamba, K. Makishima, K. Mukai, Y. Urata, A. Nitta, O. de Jager, P. Chadwick, S. Wagner, D. Buckley, M. S. Tashiro, T. Kouzu, S. Koyama, R. Yamazaki <i>X-ray searches for new galactic particle-acceleration sites with Suzaku</i>	590
E. Costantini, C.P. de Vries <i>Characterizing the chemistry of interstellar dust: the X-ray view</i>	592
A. Sartori, A. De Luca, D. Salvetti, P. Esposito, A. Tiengo, S. Zane, R. Turolla, F. Pizzolato, R. P. Mignani, P. A. Caraveo, S. Mereghetti, G. F. Bignami <i>A time-variable, phase-dependent emission line in the isolated neutron star RX J0822-4300</i>	596

	459
A. Różańska, J. Kałużny, M. Różyczka, W. Krzemiński, I.B. Thompson <i>Second neutron star in globular cluster M4</i>	598
N. Sartore, S. Mereghetti, A. Tiengo, A. De Luca, R. Turolla, F. Haberl <i>Ten years of XMM-Newton observations of RX J1856.5-3754</i>	600
P. Romano, V. Mangano, L. Ducci, P. Esposito, S. Vercellone, D.N. Burrows, J.A. Kennea, H.A. Krimm, N. Gehrels <i>From hours to years: Swift's revolutionary view of SFXTs</i>	602
V. Mangano, P. Romano, C. Ceccobello, R. Farinelli <i>IGR J08408–4503 in outburst observed by Swift</i>	604
V. Mangano, P. Esposito, P. Romano, L. Ducci, S. Vercellone <i>Swift monitoring of IGR J16418–4532</i>	606
H. Inoue <i>Precessions of accretion disks in close binaries</i>	608
A. Nebot Gómez-Morán, C. Motch <i>An XMM-Newton view of the Milky Way</i>	610
 <b>Session III: Galactic black holes and ultraluminous X-ray sources</b>	
R. Fender <i>Galactic black holes</i>	612
A. Eckart, K. Mužić, S. Yazici, N. Sabha, B. Shahzamanian, G. Witzel, L. Moser, M. Garcia-Marin, M. Valencia-S., B. Jalali, M. Bremer, C. Straubmeier, C. Rauch, R. Buchholz, D. Kunneriath, J. Moultaqa <i>The broad band spectral properties of SgrA*</i>	618
T. Enoto and Suzaku Magnetar Team <i>Broadband Suzaku observations of magnetars</i>	622
D. Radhika, A. Nandi, S. Seetha <i>Disk-Jet connection in outbursting Black Hole sources</i>	624
N. La Palombara, S. Mereghetti, L. Sidoli, A. Tiengo, P. Esposito <i>Blackbody excess in persistent Be pulsars</i>	626
J. C. Gladstone <i>The sub-classes of ultraluminous X-ray sources</i>	629
M. Mapelli <i>X-ray binaries powered by massive stellar black holes</i>	635
A. D. Sutton, T. P. Roberts, M. J. Middleton <i>Energy dependent variability as a diagnostic of the ultraluminous state: key results</i>	639
F. Pintore, P. Esposito, S. Motta, L. Zampieri <i>A stellar-mass BH in a transient, low luminosity ULX in M31?</i>	641

A. A. Nucita, F. De Paolis, R. Saxton, A. M. Read, G. Ingrosso, L. Manni <i>The high energy view of NGC 6388: hints for an IMBH?</i>	643
A. A. Nucita, L. Manni, F. De Paolis, G. Ingrosso, D. Vetrugno <i>The high energy search for IMBHs in close dSph Milky Way satellites</i>	645
<b>Session IV: X-ray background and active galactic nuclei</b>	
R. Gilli <i>The cosmic X-ray background: abundance and evolution of hidden black holes</i>	647
A. Moretti, S. Vattakunnel, P. Tozzi, R. Salvaterra, P. Severgnini, D. Fugazza, F. Haardt, R. Gilli <i>Spectrum of the unresolved cosmic X ray background</i>	653
E. Treister, C. M. Urry, K. Schawinski <i>The cosmic history of black hole growth</i>	655
A. Comastri and the XMM-CDFS Team <i>The ultra-deep XMM survey in the CDFS: X-ray spectroscopy of heavily obscured AGN</i>	661
A. Del Moro, D. M. Alexander, J. R. Mullaney, E. Daddi, F. E. Bauer, A. Pope <i>New insights on the distant AGN population</i>	665
L. W. Brenneman on behalf of the NuSTAR AGN Physics Team <i>NuSTAR's View of AGN</i>	669
A. Merloni, A. Bongiorno <i>Accreting SMBH in the COSMOS field: the connection to their host galaxies</i>	675
A. Akylas, I. Georgantopoulos, A. Georgakakis, M. Brightman, K. Nandra <i>An online tool for fitting the X-ray background and estimating the contribution of Compton-thick AGN</i>	679
P. Severgnini, A. Caccianiga, R. Della Ceca <i>The density of local Compton-thick AGN</i>	683
F. Vito, C. Vignali, R. Gilli, A. Comastri, K. Iwasawa, W. N. Brandt, D. M. Alexander, M. Brusa, B. Lehmer, F. E. Bauer, D. P. Schneider, Y. Q. Xue, B. Luo <i>The <math>z &gt; 3</math> AGN population in the 4 Ms CDFS</i>	685
M. J. Page, F. J. Carrera, J.A. Stevens, M. Symeonidis, J. D. Vieira and HerMES <i>X-ray and submillimetre observations of star-forming QSOs in the epoch of galaxy formation</i>	687
M. Cappi, F. Tombesi, M. Giustini <i>Ultra-fast outflows (aka UFOs) from AGNs and QSOs</i>	691
W. Ishibashi, T. J.-L. Courvoisier <i>The physical origin of the X-ray variability scaling in accreting black holes</i>	695

	461
A. M. Lohfink, C. S. Reynolds, R. F. Mushotzky, M. A. Nowak <i>Tackling the soft X-ray excess in AGN with variability studies</i>	699
B. De Marco <i>Soft X-ray lags and the correlation with black hole mass in radio quiet AGN</i>	703
H. Noda, K. Makishima, K. Nakazawa, S. Yamada <i>Model-independent decomposition of broad-band Suzaku spectra of AGNs into primary continua and secondary components</i>	707
A. Zoghbi <i>Fe K<math>\alpha</math> X-ray light echoes in AGN</i>	711
B. Balmaverde, A. Capetti, P. Grandi, E. Torresi, M. Chiaberge, J. Rodriguez Zaurin, G. R. Tremblay, D. J. Axon, S. A. Baum, G. Giovannini, P. Kharb, F. D. Macchetto, C. P. O'Dea, W. Sparks <i>Extended soft X-ray emission in 3CR radio galaxies at <math>z &lt; 0.3</math>: High Excitation and Broad Line Galaxies</i>	715
R. Fanali, A. Caccianiga, P. Severgnini, R. Della Ceca, M. Dotti, E. Marchese <i>The relationship between X-ray emission and accretion in X-ray selected AGNs</i>	717
<b>Session V: relativistic jets and gamma-ray bursts</b>	
G. Ghisellini <i>High redshift blazars</i>	719
M. Huarte-Espinosa, E. G. Blackman, A. Hubbard, A. Frank <i>Mass loading and knot formation in AGN jets by stellar winds</i>	725
A. Janiuk, S. Charzyński, P. Mioduszewski <i>Accretion and outflow in the gamma ray bursts from black hole binary systems</i>	727
S. Vercellone and The AGILE Team <i>The SWIFT knife: the unique SWIFT long-term monitoring program on 3C 454.3</i>	729
L. Foschini, E. Angelakis, G. Bonnoli, V. Braitto, A. Caccianiga, L. Fuhrmann, L. Gallo, G. Ghirlanda, G. Ghisellini, D. Grupe, T. Hamilton, S. Kaufmann, S. Komossa, Y. Y. Kovalev, A. Lahteenmaki, M. L. Lister, K. Mannheim, L. Maraschi, S. Mathur, B. M. Peterson, P. Romano, P. Severgnini, G. Tagliaferri, J. Tammi, F. Tavecchio, O. Tibolla, M. Tornikoski, S. Vercellone <i>Basic properties of Narrow-Line Seyfert 1 Galaxies with relativistic jets</i>	731
B. Kapanadze, S. Kapanadze <i>Swift/XRT observations of high-energy selected BL Lacertae source PKS 2155-304</i>	733
M. Ohno, Y. Hanabata, T. Kawano, K. Takaki, R. Nakamura, Y. Tanaka, Y. Fukazawa, H. Ueno, M. Tashiro, Y. Terada, W. Iwakiri, T. Yasuda, M. Asahina, S. Kobayashi, A. Sakamoto, Y. Ishida, S. Sugimoto, M. Akiyama, N. Ohmori, M. Yamauchi, Y. E. Nakagawa, S. Sugita, K. Yamaoka, M. Kokubun, T. Takahashi, Y. Urata, P. Tssai, K. Nakazawa, K. Makishima and the Suzaku-WAM team <i>All-sky observations by Suzaku wide-band all-sky monitor</i>	735


**Session VI: galaxies and clusters of galaxies**

- E. Pointecouteau  
*The Sunyaev-Zeldovich effect* 737
- F. Gastaldello, L. Di Gesu, S. Ghizzardi, S. Giacintucci, M. Girardi, E. Roediger, M. Rossetti  
*Sloshing cold fronts in the IC1860 group* 743
- P. Tozzi  
*A scientific case for future X-ray astronomy: galaxy clusters at high redshifts* 747
- Y. Wang  
*Central gas entropy excess in galaxy groups and galaxy clusters* 752
- Y. Shimoda, M. Kawaharada, K. Sato, T. Ohashi, Y. Ishisaki, I. Mitsuishi, H. Akamatsu, M. S. Tashiro  
*Metal abundances in the ICM associated with a medium-redshift cluster of galaxies MS 1512.4+3647 observed with Suzaku* 756
- I. Mitsuishi, N. Y. Yamasaki, Y. Takei  
*X-ray properties of the starburst-driven outflow in NGC 253* 758
- A. Negri, S. Pellegrini, L. Ciotti  
*Disk dynamics and the X-ray emission of S0 and flat early-type galaxies* 762
- S. Pellegrini, J. Wang, G. Fabbiano, D-W. Kim, N.J. Brassington, J.S. Gallagher, G. Trinchieri, A. Zezas  
*AGN activity and the extended hot ISM in the compact radio elliptical NGC 4278* 764
- S. Posacki, S. Pellegrini, L. Ciotti  
*The effects of flattening and rotation on the temperature of the X-ray halos of elliptical galaxies* 766
- P. Tzanavaris, T. Fragos, M. Tremmel, L. Jenkins, A. Zezas, B. D. Lehmer, A. Hornschemeier, V. Kalogera, A. Ptak, A. R. Basu-Zych  
*XSINGS: probing X-ray binaries in galaxies* 768

**Session VII: future missions and technological concepts**

- P. Predehl on behalf of the eROSITA Team  
*eROSITA* 770
- T. Takahashi  
*The ASTRO-H Mission* 776
- L. Stella (on behalf of the LOFT Team)  
*LOFT: the Large Observatory For x-ray Timing* 782
- P. Romano, P. Esposito, E. Bozzo, C. Ferrigno, V. Mangano  
*Supergiant Fast X-ray Transients with LOFT* 788

- S. S. Murray, S. Borgani, S. Campana, O. Citterio, W. Forman, R. Giacconi, R. Gilli, M. Paolillo, G. Pareschi, A. Ptak, P. Rosati, P. Tozzi, M. Weisskopf, and the WFXT Team  
*Wide Field X-ray Telescope (WFXT)* 790
- R. Petre and the NASA X-ray Mission Concepts Study Team  
*AXSIO and the NASA X-ray Mission Concepts Study* 798
- A. Vikhlinin  
*SMART-X, "Square Meter, Arcsecond Resolution Telescope for X-rays"* 805
- P. Gorenstein  
*High angular resolution X-ray astronomy in the next 50 years* 811
- M. M. Civitani, S. Campana, O. Citterio, P. Conconi, G. Pareschi, G. Tagliaferri, G. Parodi  
*High angular resolution optics for the next generation of wide field X-ray telescopes beyond e-Rosita* 817
- L. Proserpio, S. Basso, M. Civitani, O. Citterio, P. Conconi, M. Ghigo, G. Pareschi, B. Salmaso, D. Spiga, G. Tagliaferri  
*Segmented glass optics for next generation X-ray telescopes* 819
- M. Gubarev, B. Ramsey, S.O'Dell, R. Elsner, K. Kilaru, C. Atkins, D. Swartz, J. Gaskin and M. Weisskopf  
*Flight programs and X-ray optics development at MSFC* 821
- E. Virgili, F. Frontera, V. Liccardo, V. Valsan on behalf of the LAUE collaboration  
*A lens for hard X-/soft  $\gamma$ -rays: the LAUE Project* 823
- D. Götz and J. Osborne on behalf of the MXT Instrument Science Team  
*MXT: a light X-ray telescope for X-Gamma-Ray Burst afterglow observations* 825
- M. Jackson, on behalf of the PoGOLite collaboration  
*Balloon-borne hard X-ray astronomy with PoGOLite: Opening a new window on the universe* 827
- M. Barbera, A. Collura, F. Gastaldello, U. Lo Cicero, N. La Palombara, A. Tiengo, S. Varisco  
*Investigation on the status of the XMM-Newton EPIC T and M filters after more than 10 years of operation* 829
- A. Paizis, S. Mereghetti, D. Götz, M. Fiorini, M. Gaber, R. Regni Ponzeveroni, L. Sidoli, S. Vercellone  
*GOLIA: an INTEGRAL archive @INAF-IASF Milano* 831
- G. Branduardi-Raymont, S. F. Sembay, J. P. Eastwood, D. G. Sibeck, J. A. Carter, A. M. Read and the AXIOM Collaboration  
*X-ray imaging of the Earth's magnetosphere* 833
- C. M. Urry  
*Planning for future X-ray astronomy missions* 835



# The effects of flattening and rotation on the temperature of the X-ray halos of elliptical galaxies

S. Posacki, S. Pellegrini, and L. Ciotti

Department of Physics and Astronomy, University of Bologna, Viale Berti Pichat 6/2, I-40127 Bologna, Italy, e-mail: [silvia.posacki@unibo.it](mailto:silvia.posacki@unibo.it)

**Abstract.** Elliptical galaxies have hot coronae with X-ray luminosities and mean gas temperatures that span over wide ranges. This variation can be partially due to the energy budget of the hot gas, that depends on the host galaxy structure and internal kinematics. With the aid of realistic axisymmetric galaxy models, we performed a diagnostic study focussed on the effects of galaxy flattening and rotational support on the hot gas temperature.

**Key words.** Galaxies: elliptical and lenticular, cD – Galaxies: fundamental parameters – Galaxies: ISM – Galaxies: kinematics and dynamics – X-rays: galaxies – X-rays: ISM

## 1. Introduction


The X-ray observatory *Chandra* allowed for a study of the hot gas halos in early-type galaxies (ETGs) with unprecedented quality and accuracy. With these data, an ETG sample with homogeneous measurements of the average hot gas temperature and luminosity has recently been built (Boroson et al., 2011), and provides the basis to test the hot gas properties against other major galaxy properties. For example, the hot gas content has long been known to be sensitive to the shape of the mass distribution, and possibly to the mean rotation velocity of the stars (see Pellegrini 2012 for a recent review). The same could be concluded in a recent study focussed on the gas temperature (Pellegrini, 2011). Here we show some preliminary results from an investigation of the effects of galaxy shape and internal kinematics on the gas temperature, and on the energy re-

quired to extract the gas from the galaxy potential well, for axisymmetric ETG models.

## 2. The galaxy models

The galaxy models include three mass components: a stellar distribution following the approximate ellipsoidal deprojection of the de Vaucouleurs profile, with axial ratio  $0.3 \leq q \leq 1$ ; a spherically symmetric dark matter (DM) halo, described either by the singular isothermal sphere (SIS), the Hernquist, the Einasto, or the Navarro-Frenk-White profile; and a central massive black hole. We assume a two-integral phase-space distribution function, so that  $\sigma_R = \sigma_z$ , and the amount of ordered rotational support in the azimuthal direction is controlled by the Satoh (1980)  $k$ -decomposition:  $k = 1$  corresponds to the isotropic rotator, whereas for  $k = 0$  no net rotation is present, and the galaxy flattening is due to  $\sigma_\varphi$  only. The Jeans equations are solved with





**Fig. 1.**  $T_*$  (top) and  $T_{\text{grav}}$  (bottom) for two EO-built sub-families with  $\sigma_{e8} = 150$  and  $300 \text{ km s}^{-1}$ , as a function of morphological type and  $(k, \gamma)$ . Round and elliptical symbols refer to the FO and EO view of the same models, respectively. Filled and empty symbols are for  $k = 0$  and  $k = 1$ .  $\gamma = (1, 0.5, 0)$  decreases from top to bottom for the empty symbols. The shaded area shows the range spanned by  $T_{\text{inj}}$ .

a numerical code built on purpose, under the assumption of a constant stellar mass-to-light ratio. All the structural and dynamical properties are projected face-on (FO) and edge-on (EO), and the circularized effective radius ( $R_e$ ) and the luminosity weighted velocity dispersion within  $R_e/8$  ( $\sigma_{e8}$ ) are computed. The models are forced to follow the most important scaling laws, so that  $(R_e, \sigma_{e8}, L)$  satisfy the Faber-Jackson and Size-Luminosity relations. Finally we calculate fiducial ISM temperatures for the models, as defined in Pellegrini (2011), under the assumption that the mass sources are distributed as the stellar density. The temperature  $T_* = T_\sigma + \gamma T_{\text{rot}}$  measures the energy input due to the thermalization of stellar random ( $T_\sigma$ ) and ordered ( $T_{\text{rot}}$ ) motions. The actual value of  $\gamma$  is not known a priori, since it depends on the relative motion between the freshly injected gas and the gas already permeating the galaxy. It can be obtained only from hydrodynamical simulations, that suggest it assumes small values (see Negri et al., this volume). The injection temperature  $T_{\text{inj}} = T_* + T_{\text{SN}}$  takes into account also the contribution of

the kinetic energy of SNIa explosions ( $T_{\text{SN}}$ ). Finally,  $T_{\text{grav}}$  quantifies the average energy per unit mass required to extract the gas from the galaxy potential well.

### 3. Results

We explored how  $T_*$ ,  $T_{\text{inj}}$  and  $T_{\text{grav}}$  depend on  $(q, k, \gamma)$ . The results presented here refer to a SIS halo truncated at  $15 R_e$ . The model families are built starting from a spherical object with luminosity  $L$  and effective radius  $R_e$  that represents the progenitor of a given family. The flattened models belonging to a family are built varying  $q$ , while keeping both  $(L, R_e)$  and the DM halo fixed. Being  $R_e$  dependent on the line-of-sight (l.o.s.) direction, each family consists of two sub-families: a FO and an EO-built one. As a result,  $\sigma_{e8}$  and the temperatures depend on the choice of  $(q, k, \gamma)$  and l.o.s. direction. Figure 1 shows  $T_*$ ,  $T_{\text{grav}}$  and  $T_{\text{inj}}$  for two EO-built sub-families. The models reproduce the observed correlation between the mean ISM temperature and  $\sigma_{e8}$  (e.g. Boroson et al. 2011), an aspect discussed in Posacki et al. (in preparation). The more the galaxy is flattened, the larger can be the effect of rotation: flatter models allow more rotational support, so that  $T_*$  spans wider ranges as a function of  $k$  (and consequently of  $\gamma$ ). Instead,  $k$  variations have only mild effects on  $T_*$  of rounder systems. Both  $T_*$  and  $T_{\text{grav}}$  slightly decrease with  $q$ , and the  $T_{\text{inj}}$  values are dominated by  $T_{\text{SN}}$ . If  $T_{\text{inj}} > T_{\text{grav}}$  the gas can escape from the galaxy (see Pellegrini 2011 for a detailed discussion), thus smaller galaxies are more likely to host an outflow/wind region than bigger ones, as already observed (e.g. Pellegrini 2012). We found that these temperature trends are almost independent of the specific DM halo profile.

### References

- Boroson, B., Kim, D.-W., & Fabbiano, G. 2011, ApJ, 729, 12
- Pellegrini, S. 2011, ApJ, 738, 57
- Pellegrini, S. 2012, , Astrophysics and Space Science Library, 378, 21
- Sato, C. 1980, PASJ, 32, 41