

IAU Symposium

315

3–7 August 2015  
Honolulu, USA

Proceedings of the International Astronomical Union

# From Interstellar Clouds to Star-forming Galaxies: Universal Processes?

*Edited by*


Pascale Jablonka

Philippe André

Floris van der Tak

ISSN 1743-9213

International Astronomical Union



CAMBRIDGE  
UNIVERSITY PRESS



INTERNATIONAL ASTRONOMICAL UNION  
UNION ASTRONOMIQUE INTERNATIONALE

International Astronomical Union



FROM INTERSTELLAR CLOUDS  
TO STAR-FORMING GALAXIES:  
UNIVERSAL PROCESSES?

PROCEEDINGS OF THE 315th SYMPOSIUM  
OF THE INTERNATIONAL ASTRONOMICAL  
UNION HELD IN HONOLULU, HAWAII,  
UNITED STATES  
AUGUST 3–7, 2015

Edited by

PASCALE JABLONKA

*EPFL, Switzerland*

PHILIPPE ANDRÉ

*CEA - AIM Paris-Saclay, France*

and

FLORIS VAN DER TAK

*SRON & University of Groningen, The Netherlands*



CAMBRIDGE  
UNIVERSITY PRESS

CAMBRIDGE UNIVERSITY PRESS  
University Printing House, Cambridge CB2 8BS, United Kingdom  
32 Avenue of the Americas, New York, NY 10013, USA  
10 Stamford Road, Oakleigh, Melbourne 3166, Australia

© International Astronomical Union 2016

This book is in copyright. Subject to statutory exception  
and to the provisions of relevant collective licensing agreements,  
no reproduction of any part may take place without  
the written permission of the International Astronomical Union.

First published 2016

Printed in the UK by Bell & Bain, Glasgow, UK

Typeset in System L<sup>A</sup>T<sub>E</sub>X 2<sub>ε</sub>

*A catalogue record for this book is available from the British Library Library of Congress  
Cataloguing in Publication data*

This journal issue has been printed on FSC<sup>TM</sup>-certified paper and cover board. FSC is an  
independent, non-governmental, not-for-profit organization established to promote the  
responsible management of the world's forests. Please see [www.fsc.org](http://www.fsc.org) for information.

ISBN 9781107135208 hardback  
ISSN 1743-9213

PASCAL JABLONKA  
EPFL, Switzerland  
PHILIPPE ANDRÉ  
CEA - AIM Paris-Saclay, France

and  
FLORIS VAN DER TAK

CAMBRIDGE  
UNIVERSITY PRESS



# AGN feedback and star formation in ETGs: negative and positive feedback

Luca Ciotti<sup>1</sup>, Jeremiah P. Ostriker<sup>2,3</sup>, Andrea Negri<sup>1,4</sup>,  
Silvia Pellegrini<sup>1</sup>, Silvia Posacki<sup>1</sup> and Greg Novak<sup>5</sup>

<sup>1</sup>Dept. of Physics and Astronomy, University of Bologna, via Ranzani 1, I-40127 Bologna, Italy,  
email: luca.ciotti@unibo.it

<sup>2</sup>Dept. of Astrophysical Sciences, Princeton University, NJ, USA

<sup>3</sup>Dept. of Astronomy and Astrophysics, Columbia University, NY, USA

<sup>4</sup> Institut d'Astrophysique de Paris, CNRS, UMR 7095 98bis bvd Arago, F-75014 Paris, France

<sup>5</sup> Observatoire de Paris, LERMA, CNRS, 61 Av de l'Observatoire, F-75014 Paris, France

**Abstract.** AGN feedback from supermassive black holes (SMBHs) at the center of early type galaxies is commonly invoked as the explanation for the quenching of star formation in these systems. The situation is complicated by the significant amount of mass injected in the galaxy by the evolving stellar population over cosmological times. In absence of feedback, this mass would lead to unobserved galactic cooling flows, and to SMBHs two orders of magnitude more massive than observed. By using high-resolution 2D hydrodynamical simulations with radiative transport and star formation in state-of-the-art galaxy models, we show how the intermittent AGN feedback is highly structured on spatial and temporal scales, and how its effects are not only negative (shutting down the recurrent cooling episodes of the ISM), but also positive, inducing star formation in the inner regions of the host galaxy.

**Keywords.** ISM: kinematics and dynamics, galaxies: active, galaxies: cooling flows, galaxies: elliptical and lenticular, cD, galaxies: starburst, galaxies: evolution

## 1. Introduction

In a quite widespread view, merging is considered the main cause of QSO activity, because it is believed to be, after the end of the galaxy formation epoch, the only mechanism to add fresh gas to the central SMBHs in early type galaxies (ETGs). However, it has also been shown (e.g., Ciotti & Ostriker 1997) that the stellar mass losses produced during stellar evolution cyclically feed a central gas inflow, and then trigger the QSO activity in isolated ETGs (sometimes considered “red and dead”). Indeed, recent observations indicate that the QSO activity can be independent of galaxy merging. QSO activity is also invoked as the explanation of star formation quenching in ETGs (negative feedback) but, as stressed in Ciotti & Ostriker (2007, hereafter CO07), AGN feedback can induce star formation (positive feedback) during “cooling flow” episodes fueled by stellar mass losses. In fact, also in isolation, significant amounts of fresh gas ( $\simeq 20 - 30\%$  of the initial stellar mass  $M_*$  of the galaxy, depending on the IMF) are injected over the galaxy body by stellar evolution, at a rate approximately given by  $\dot{M}_*(t) \propto M_*(0) t^{-1.3}$  (Ciotti *et al.* 1991, see also Pellegrini 2012). These losses interact with the pre-existing hot ISM, and mix with it, due to thermalization of the stellar velocity dispersion. In a galaxy of total B-band luminosity  $L_B$ , SNIa's explosions at a rate  $R_{\text{SN}}(t) \propto L_B t^{-s}$  provide additional mass and heat to the ISM, with rates  $\dot{M}_{\text{SN}}(t) = 1.4 M_\odot R_{\text{SN}}(t)$  and  $L_{\text{SN}}(t) = 10^{51} \text{ erg } R_{\text{SN}}(t)$ . Since recent estimates suggest  $s \simeq 1$ , then the SNIa's specific heating  $L_{\text{SN}}/\dot{M}_*$  increases with time. The resulting hot atmosphere however cannot be


static. The interstellar mass keeps increasing, its cooling time decreases, until a cooling catastrophe takes place in the central galactic region and the SMBH feedback takes place.

In a series of papers, Ciotti, Ostriker and co-workers, with the aid of high-resolution hydrodynamical simulations, and a physically based feedback implementation, studied in detail the AGN activity cycle induced by accretion of stellar mass losses, also considering star formation. In particular, it was shown that AGN feedback is able to avoid the overgrowth of the SMBH that would be caused by accretion of the mass lost by stars if unimpeded, and also that the same feedback can induce significant star formation at the center of ETGs (CO07). By using state-of-the-art galaxy dynamical models, and an updated version of the 2D code of Novak *et al.* (2011, 2012, hereafter N12, and references therein), we are now investigating in detail the ISM behavior in realistic ETG models with evolving input ingredients from the stellar population (Ciotti *et al.*, in preparation).

## 2. The models

We consider axisymmetric two-component galaxy models with adjustable flattening embedded in a spherically symmetric NFW dark matter halo and with a central SMBH. The stellar component is described by the ellipsoidal deprojection of the de Vaucouleurs law. The total gravitational field, the solution of the two-integrals Jeans equations, and the projection of the resulting kinematical field on the plane of the sky are computed as described elsewhere (Posacki *et al.* 2013). The velocity dispersion and rotational fields of the stellar populations, needed to compute energy and momentum injection in the ISM due to stellar evolution, are computed by using a generalized Satoh  $k$ -decomposition (Ciotti & Pellegrini 1996). The E4 and E7 galaxy models are obtained by flattening (at fixed stellar mass  $M_*$ ) spherical “progenitors” with central (aperture) velocity dispersion of 180, 210, 250 and 300 km/s, and reproduce observed scaling laws (Negri *et al.* 2014). The initial mass of the SMBH is taken to be  $M_{\text{BH}} = 10^{-3} M_*$ , near to the Magorrian relation, as expected at the end of the period of galaxy formation.

The hydrodynamical equations and the input physics are given in Ciotti & Ostriker (2012), and integrated by an improved version of the N12 code. Mass, momentum and energy sources and sinks associated with evolution of the passively evolving initial stellar population, and with the new stars added by star formation, are treated as described in Negri *et al.* (2015, hereafter N15). In particular the star formation rate is given by  $\dot{\rho}_{\text{SF}} = 0.1\rho/\max(\tau_{\text{cool}}, \tau_{\text{dyn}})$ , where  $\rho$  is the local ISM density, and  $\tau_{\text{cool}}$  and  $\tau_{\text{dyn}} = \min(\tau_{\text{Jeans}}, \tau_{\text{rot}})$  are the cooling and dynamical times, respectively. In N15 we showed that this prescription reproduces the observed Kennicutt-Schmidt law remarkably well. Radiative and mechanical feedback from the accreting SMBH (both depending on the hydrodynamically self-consistently determined mass accretion rate on the SMBH,  $\dot{M}_{\text{BH}}$ ), are finally considered. In the energy equation, photoionization and Compton heating and cooling ( $T_{\text{C}} \simeq 2 \times 10^7$  K), bremsstrahlung and line cooling are taken into account, with modifications due to the ionization effects of AGN radiation. In the momentum equation, radiation pressure due to AGN activity is computed by solving the radiative transport (in spherical symmetry) for the accretion luminosity  $L_{\text{BH}}$ , considering photoionization+Compton opacity, and electron scattering. For simplicity at this stage we ignore the detailed treatment of dust formation and destruction, and the associated radiation pressure effects (Hansley *et al.* 2014, N12), as well as the radiation pressure due to star light in different bands (e.g., see Ciotti & Ostriker 2012). For given  $\dot{M}_{\text{BH}}$ , then,



**Figure 1.** Global properties of the models at the end of the simulations (13 Gyr), as a function of their initial stellar mass, for E4 (black) and E7 (red) models. Triangles, squares, and pentagons refer to models with radiative plus mechanical AGN feedback, mechanical AGN feedback only, and no AGN feedback, respectively. *A color version of this figure is available online.*

one has:

$$L_{\text{BH}} = \epsilon_{\text{EM}} \dot{M}_{\text{BH}} c^2, \quad \epsilon_{\text{EM}} = \frac{\epsilon_0 A \dot{m}}{1 + A \dot{m}}, \quad \dot{m} \equiv \frac{\dot{M}_{\text{BH}}}{\dot{M}_{\text{Edd}}} = \frac{\epsilon_0 \dot{M}_{\text{BH}} c^2}{L_{\text{Edd}}}, \quad (2.1)$$

where  $A = 100$ ,  $\epsilon_0 = 0.125$  and  $L_{\text{Edd}}$  is the Eddington luminosity.

Mechanical feedback is due to mass, momentum, and energy injection of a conical-shaped nuclear wind, with half-opening angle of  $\simeq 45^\circ$ . In terms of solid angle, this means that the wind is visible from  $\sim 1/4$  of the available viewing angles. The relevant relations here are  $\dot{M}_{\text{BH}} = \dot{M}_{\text{in}}/(1 + \eta)$ ,  $\dot{M}_{\text{out}} = \eta \dot{M}_{\text{BH}}$ ,  $\dot{p}_{\text{w}} = \dot{M}_{\text{out}} v_{\text{w}}$ ,  $L_{\text{w}} = \epsilon_{\text{w}} \dot{M}_{\text{BH}} c^2$ , where  $\eta \equiv 2\epsilon_{\text{w}} c^2/v_{\text{w}}^2$ ,  $\dot{M}_{\text{in}}$  and  $\dot{M}_{\text{out}}$  are the mass inflow and outflow rates at the first grid point (a few pc from the SMBH). In the current simulations,  $\epsilon_{\text{w}}$  follows a prescription similar to that adopted for  $\epsilon_{\text{EM}}$ , with maximum value of  $10^{-4}$ , while the wind velocity  $v_{\text{w}}$  is  $10^4$  km/s.

### 3. Results

As illustrated in Fig. 1 (top panel), the central SMBH grows more, in absolute terms ( $\Delta M_{\text{BH}}$ ) and also in percentage ( $\Delta M_{\text{BH}}/M_{\text{BH}}(0)$ ), for increasing stellar mass  $M_*$  of the host galaxy; the growth reaches a factor of  $\simeq 3$  in the biggest galaxies with AGN feedback (to be compared with a factor  $\simeq 30$  in models without AGN feedback). Moreover, at

any fixed  $M_*$ , more mass is accreted by the SMBH in E4 galaxies than in E7 galaxies. This is due to the fact that edge-on flattened galaxies are less bound (see N15) than more spherical systems. As expected, at fixed  $M_*$  and galaxy shape, the accreted mass increases from models with full AGN feedback (triangles), to models with mechanical AGN feedback only (squares), and finally to models without AGN feedback (pentagons). Therefore, since we start with a SMBH mass  $\sim$  on the Magorrian relation, this relation is quite well preserved thanks to the effect of AGN feedback, which is able, in combination with large-scale SNIa heating, to maintain small the SMBH masses, avoiding the accretion of  $\simeq 99\%$  of the mass injected by the stars over the galaxy body.

The stellar mass added to the galaxy due to star formation in the gas supplied by evolving stars ( $\Delta M_*$ , middle panel), is larger in E4 galaxies than in E7 galaxies of same  $M_*$ . The total mass in new stars is of the order of 4 – 5% of the initial  $M_*$ , indicating that an important fraction of the injected mass ( $\simeq 0.2 - 0.3M_*$ ) escapes as a galactic wind, supported by SNIa heating with the additional contributions of the thermalization of stellar winds, of type SNII explosions in the new stellar populations, and of the AGN feedback. In general, as already found in the case of spherical models (CO07), models with AGN feedback tend to form *more* stars than models without (with the exception of very low mass galaxies, where AGN feedback is able to sustain global galactic winds). This positive feedback is explained by considering that the AGN prevents the rapid fall of the cooling material to the center, allowing for more time for star formation. A clear sign of this effect is the characteristic size of the region of star formation, which is confined to the very center of the galaxy in absence of AGN feedback (a small fraction of the optical effective radius), while in models with AGN feedback it reaches a size of the order of a kpc or more. The results of this work (in absence of AGN feedback) agree well with those extensively discussed in N15 including star formation.

An important observational feature of the models is of course the evolution of their luminosity, both of the central accreting SMBH ( $L_{\text{BH}}$ ), and of the galaxy hot ISM ( $L_X$ ). In Fig. 1 we show  $L_X$  in the X-ray 0.3 – 8 keV band at the end of the simulations, with values in nice agreement with observations, and only moderately dependent on AGN feedback. The evolution of  $L_{\text{BH}}$  (not shown), is confirmed to be highly fluctuating, as already found in our previous works based on simpler galaxy models. The duty-cycle (phenomenologically defined as the fraction of time spent by the SMBH at  $L_{\text{BH}}$  higher than a few percents of  $L_{\text{Edd}}$ ), is of the order of  $\simeq 3 - 5\%$ .

## References

- Ciotti, L., D’Ercole, A., Pellegrini, S., & Renzini, A. 1991, *ApJ*, 376, 380  
 Ciotti, L. & Ostriker, J. P. 1997, *ApJ*, 487, L105  
 Ciotti, L. & Ostriker, J. P. 2007, *ApJ*, 665, 1038 (CO07)  
 Ciotti, L. & Ostriker, J. P. 2012, in D.W. Kim and S. Pellegrini (eds.), *Hot interstellar matter in elliptical galaxies*, (Springer, Heidelberg), ASSL, vol. 378, p. 83  
 Ciotti, L. & Pellegrini, S. 1996, *MNRAS*, 279, 240  
 Hensley, B., Ostriker, J. P., & Ciotti, L. 2014, *ApJ*, 789, 78  
 Negri, A., Posacki, S., Pellegrini, S., & Ciotti, L. 2014, *MNRAS*, 445, 1351  
 Negri, A., Pellegrini, S., & Ciotti, L. 2015, *MNRAS*, 451, 1212 (N15)  
 Novak, G., Ostriker, J. P., & Ciotti, L. 2011, *ApJ*, 737, 26  
 Novak, G., Ostriker, J. P., & Ciotti, L. 2012, *MNRAS*, 427, 2734 (N12)  
 Pellegrini, S. 2012, in D.W. Kim and S. Pellegrini (eds.), *Hot interstellar matter in elliptical galaxies*, (Springer, Heidelberg), ASSL, vol. 378, p. 21  
 Posacki, S., Pellegrini, S., & Ciotti, L. 2013, *MNRAS*, 433, 2259

# Star formation in early-type galaxies: the role of stellar winds and kinematics

Silvia Pellegrini<sup>1</sup>, Andrea Negri<sup>2</sup> and Luca Ciotti<sup>1</sup>

<sup>1</sup>Dept. of Physics and Astronomy, University of Bologna, via Ranzani 1, I-40127 Bologna, Italy  
email: [silvia.pellegrini@unibo.it](mailto:silvia.pellegrini@unibo.it)

<sup>2</sup>CNRS, UMR 7095, Institut d'Astrophysique de Paris, 98bis bvd Arago, F-75014 Paris, France

**Abstract.** Early-type galaxies (ETGs) host a hot ISM produced mainly by stellar winds, and heated by Type Ia supernovae and the thermalization of stellar motions. High resolution 2D hydrodynamical simulations showed that ordered rotation in the stellar component results in the formation of a centrifugally supported cold equatorial disc. In a recent numerical investigation we found that subsequent generations of stars are formed in this cold disc; this process consumes most of the cold gas, leaving at the present epoch cold masses comparable to those observed. Most of the new stellar mass formed a few Gyrs ago, and resides in a disc.

**Keywords.** stars: formation; ISM: evolution; ISM: kinematics and dynamics; galaxies: elliptical and lenticular, cD; galaxies: ISM; galaxies: structure

---


## 1. Introduction

High resolution 2D hydrodynamical simulations with the ZEUS-MP2 code showed that ordered rotation in the stellar component affects significantly the evolution of the hot ISM in ETGs, and, among other effects, results in the formation of a centrifugally supported cold equatorial disc (Negri *et al.* 2014, hereafter N14). This disc can be extended ( $\sim 0.5 - 10$  kpc radius), and as massive as  $10^{10} M_{\odot}$  in the biggest ETGs. Indeed there is evidence that  $\sim 50\%$  of massive ETGs host significant quantities of cold gas (Davis *et al.* 2011, Serra *et al.* 2014), often in settled configurations, sharing the same kinematics of the stars, consistent with an internal origin. Also, the cold gas seems to provide material for low level star formation (hereafter SF); and, in the ATLAS<sup>3D</sup> sample, molecular gas, SF and young stellar populations are detected only in fast rotators (Sarzi *et al.* 2013, Davis *et al.* 2014). We then added the possibility for the gas to form stars to the simulations of N14, to explore whether SF can bring the amount of cold gas in the models more in agreement with observed values, and whether it can explain the low-level SF activity currently seen to be ongoing in rotating systems.

## 2. The models and the results

Hydrodynamical simulations were run for a representative subset of 12 rotating models from N14, including in the code the removal of cold gas, and the injection of mass, momentum and energy appropriate for the newly forming stellar population (Negri *et al.* 2015). SF was implemented by subtracting gas from the grid, at an adopted rate per unit volume of  $\dot{\rho}_{\text{SF}} = \eta_{\text{SF}} \rho / t_{\text{SF}}$ , where  $\rho$  is the gas density,  $\eta_{\text{SF}}$  is the SF efficiency ( $\eta_{\text{SF}} = 0.01$  and  $0.1$  were adopted), and  $t_{\text{SF}}$  is the maximum between the cooling timescale and the dynamical timescale. In a typical (cyclical) evolution, gas injected by the stellar population accumulates until radiative losses become catastrophic, significant amounts of cold material are produced, and SF is enhanced. At the end the cold gas mass  $M_c$  is much reduced in the models with SF, and the mass in new stars  $M_{\text{new}}^*$  is close to the  $M_c$  values of the models without SF (Fig. 1). The new stars reside mostly in a disc, and could be related to a younger, more metal rich disk-like stellar component observed in fast rotators (Cappellari *et al.* 2013). Most of  $M_{\text{new}}^*$  formed a few Gyrs ago; the SF rate at the present epoch is low ( $\leq 0.1 M_{\odot} \text{yr}^{-1}$ ), as observed, at least for model ETGs of stellar





**Figure 1.** From top to bottom, final values for all rotating models of: the cold gas mass  $M_c$  [without SF (left panel), and with the two adopted  $\eta_{\text{SF}}$  values (other panels)]; the stellar mass in newly formed stars  $M_*^{\text{new}}$ ; the SF rate normalized to the initial stellar mass of the galaxy  $M_*$ ; the mean formation time of the new stars  $\langle t \rangle_*^{\text{new}}$ , calculated from an initial time of 2 Gyr. See Negri *et al.* (2015) for more details.

mass  $< 10^{11} M_\odot$  (Fig. 1). The adopted SF recipe reproduces the slope of the Kennicutt-Schmidt relation, and even its normalization for  $\eta_{\text{SF}} = 0.01$  (Negri *et al.* 2015).

## References

- Cappellari, M., *et al.* 2013, *MNRAS*, 432, 1862  
 Davis, T.A., *et al.* 2011, *MNRAS*, 417, 882  
 Davis, T. A., *et al.* 2014, *MNRAS*, 444, 3427  
 Negri, A., Posacki, S., Pellegrini, S., & Ciotti, L. 2014, *MNRAS*, 445, 1351 (N14)  
 Negri, A., Pellegrini, S., & Ciotti, L. 2015, *MNRAS*, 451, 1212  
 Sarzi, M., *et al.* 2013, *MNRAS*, 432, 1845  
 Serra, P., *et al.* 2014, *MNRAS*, 444, 3388