

ORIGINAL ARTICLE

Metal abundances in the MACER simulations of the hot interstellar medium

Silvia Pellegrini¹ | Zhaoming Gan^{2,3} | Jeremiah P. Ostriker³ | Luca Ciotti¹

¹Department of Physics and Astronomy,
University of Bologna, Bologna, Italy

²Shanghai Astronomical Observatory,
Chinese Academy of Sciences, Shanghai,
People's Republic of China

³Department of Astronomy, Columbia
University, New York, New York

Correspondence

Silvia Pellegrini, Department of Physics
and Astronomy, University of Bologna, via
Gobetti 93/2, I-40129 Bologna, Italy.
Email: silvia.pellegrini@unibo.it

Abstract

Hot plasma is the dominant phase of the interstellar medium of early-type galaxies. It can originate from stellar mass losses, residual gas from the formation epoch, and accretion from outside the galaxies. Its evolution is linked to the dynamical structure of the host galaxy, to the supernova and active galactic nuclei (AGN) feedback, and to (late-epoch) star formation in a way that has yet to be fully understood. Important clues about the origin and evolution of the hot gas come from the abundances of heavy metals that have been studied with increasing detail through *XMM-Newton* and *Chandra*. We present recent high-resolution hydrodynamical simulations of the hot gas evolution that include the above processes, where several chemical species, originating in asymptotic giant branch stars and supernovae of types Ia and II, have also been considered. The high resolution, of few parsecs in the central galactic region, allows us to track the metal enrichment, transportation, and dilution throughout the galaxy. The comparison of model results with observed abundances demonstrates good agreement for the region enriched by the AGN wind but also shows discrepancies for the diffuse hot gas; the latter indicates the need for a revision of standard assumptions and/or the importance of neglected effects due to the dust and/or residual uncertainties in deriving abundances from the X-ray spectra.

KEYWORD

galaxies: active – galaxies: ISM – ISM: abundances – ISM: evolution; X-rays: ISM

1 | INTRODUCTION

Hot plasma dominates the interstellar medium (ISM) of early-type galaxies, as discovered using *Einstein* X-ray observations, and were investigated thoroughly until recently through *Chandra* and *XMM-Newton* (Fabbiano 2012). The X-ray emission of this hot phase correlates with the main galaxy properties, such as the optical luminosity and the stellar and total galaxy mass, although with a significant scatter (Forbes et al. 2017; Goulding et al.

2016). The observed X-ray properties have led to a picture where the origin of the hot ISM is in the old, aging stellar population, and perhaps also in the circumgalactic infall (Pellegrini et al. 2018; Werner et al. 2019); its evolution is the result of the main galactic properties, such as the depth and shape of the potential well, and the stellar kinematics and of important phenomena, such as heating from supernovae and active galactic nuclei (AGN) feedback and late-epoch star formation (Ciotti et al. 2017; Gan et al. 2019b; Negri et al. 2014). While the respective roles of these

processes are not fully understood yet, they are expected to contribute to the enrichment of the gas in metals, and during its evolution, they are expected to be able to circulate the metals within and outside the galaxies. Thus, metals provide important diagnostic tests for the ISM origin and the mentioned processes regulating its evolution. In the hydrodynamical MACER code for the evolution of the ISM of massive elliptical galaxies (Gan et al. 2019b), element tracers have recently been added in order to track the metal enrichment, transportation, and dilution throughout the galaxy and outside (Gan et al. 2019a). In the following, we briefly describe the MACER modeling (Section 2), with particular emphasis on the sources of mass and metals for the hot ISM (Section 2.1); we then present the results of the simulations for the metals only (Section 3) and the conclusions (Section 4), which include a comparison with observations.

2 | MODELING THE ISM EVOLUTION

Physical modeling of the ISM evolution has been implemented in grid-type hydrodynamical simulations, performed with the Athena++ code (Gan et al. 2019a,b). The main advantages of these simulations are the accurate treatment of the sources of mass and heating, the latter from SNe and the AGN, and the high resolution, which reaches 2.5 pc at the center, so that the fiducial Bondi radius for accretion onto the central massive black hole (MBH) is resolved. The ISM evolution is followed from an age of 2 Gyr (after the main galaxy formation phase) until the present epoch, for a time interval of 11 Gyr.

The code solves the time-dependent Eulerian hydrodynamical equations with sources and sinks of mass, energy, and momentum. Sources of mass are provided by three channels: (a) the original, aging stellar population; (b) the new stars (cold, dense gas is allowed to form stars, following both a Jeans criterion and Toomre instability criterion; see Gan et al. 2019b for more details); and (c) accretion from the circumgalactic medium (CGM). No mass loading of cold gas into the hot phase is considered. These mass sources, in turn, provide metals (Section 2.1). Sources of heating for the gas come from SNe explosions (of types SNIa and SNII, the latter only from the new stars, with a thermalization efficiency of 0.85), from the thermalization of the stellar motions, and from accretion onto the MBH. Accretion produces a nuclear luminosity, with an efficiency scaling with the mass accretion rate, with the consequent radiative feedback; it also produces mechanical feedback via AGN winds, with maximum velocities of 10^4 km s^{-1} and a mechanical efficiency again scaling with the mass accretion rate, as observed for nuclear outflows

(Arav et al. 2013; Carniani et al. 2015; Tombesi et al. 2015). More details are given in Gan et al. 2019b.

An important feature of the modeling is the implementation of streaming velocity for the stars; this imparts net angular momentum to their mass losses so that the gas, when infalling in a cooling flow, settles in a rotationally supported disk-like structure; gas then cools in the disk, where it becomes gravitationally unstable, according to the Toomre Instability, and new stars form. Angular momentum transfer and star formation then occur at the same time, with the consequent near coincidence of star formation and AGN activity.

2.1 | Sources of mass and metals

Mass sources to the ISM are due to stars and the CGM. The initial, old stellar population (SSP) injects mass via asymptotic giant branch (AGB) winds, at a mass rate \dot{M}_\star and, to a much lesser extent, via SNIa's ejecta at a mass rate \dot{M}_{Ia} . As described in Pellegrini (2012), the rates of these processes, after an age $t_{\text{age}} = 2 \text{ Gyr}$ for the SSP, are:

$$\dot{M}_\star = 10^{-12} A \frac{M_\star}{M_\odot} \left(\frac{t_{\text{age}}}{12 \text{ Gyr}} \right)^{-1.3} M_\odot \text{ yr}^{-1} \quad (1)$$

where $A = 3.3$ for the Kroupa IMF, and M_\star is the SSP mass at $t_{\text{age}} = 12 \text{ Gyr}$, and:

$$\dot{M}_{\text{Ia}} = 0.216 \times 10^{-12} \frac{L_B}{L_{B\odot}} \left(\frac{t_{\text{age}}}{12 \text{ Gyr}} \right)^{-1.1} M_\odot \text{ yr}^{-1} \quad (2)$$

where L_B is the present-epoch B-band luminosity of the SSP, and $1.35 M_\odot$ for the mass ejected in one SNIa event and a Hubble constant of $70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ have been assumed.

New stars form in the cold gas disk with a top-heavy IMF of an index equal to -1.65 based on observations (e.g., the central stellar disks in the Milky Way and M31) and theoretical expectations (Bartko et al. 2010; Jiang & Goodman 2011; Lu et al. 2013). The chosen IMF has $\sim 60\%$ of the new star mass in massive stars, which turn into SNII; for the new stars, only SNII provides a source of mass and metals to the ISM (Gan et al. 2019a).

Infall of the CGM onto the galaxy also provides mass to the ISM at a rate (\dot{M}_{CGM}) estimated from cosmological zoom-in simulations based on the evolution of the accretion rate for 30 massive ellipticals (Brennan et al. 2018, Gan et al. 2019b). The rate declines sharply as expected in a Λ -CDM universe. Figure 1 shows a comparison of the mass input rates from the SSP and the CGM infall during the time span of the simulations. We adopt two




FIGURE 1 The mass input rates to the ISM from the old stellar population (AGB winds, in black, \dot{M}_* ; SNIa's ejecta, in red, \dot{M}_{Ia}) and from the cosmological CGM infall (in blue, \dot{M}_{CGM} , for the two normalizations of CGM₂₀ and CGM₇ models). The \dot{M}_* and \dot{M}_{Ia} rates refer to the galaxy with $M_* = 3.3 \times 10^{11} M_\odot$ of Section 3. See Section 2.1 for more details. ISM, interstellar medium.

normalizations for the CGM infall rate, within the range shown by cosmological simulations; the corresponding total accreted mass during the time interval in Figure 1 is 1/20 (for CGM₂₀ models) and 1/7 (for CGM₇ models) of M_* . For comparison, the total injected mass from the SSP amounts to $M_*/10$ during the same time.

The three sources of mass considered above (old and new stars and the CGM) also provide metals to the ISM. In the code, the chemical evolution of the ISM due to the injection, dilution, and transportation of metals is followed, introducing 12 tracers X_i ($i = 1, 2, \dots, 12$), where X_i is the mass of the i -th element per unit volume. The elements considered are H, He, C, N, O, Ne, Mg, Si, S, Ca, Fe, and Ni. Thus, 12 additional continuity equations are solved, assuming the chemical species comove after they are injected into the ISM:

$$\frac{\partial X_i}{\partial t} + \nabla \cdot (X_i \mathbf{v}) + \nabla \cdot \dot{\mathbf{m}}_{Q,i} = \dot{X}_{i,*} + \dot{X}_{i,Ia} + \dot{X}_{i,II} - \dot{X}_{i,SF}, \quad (3)$$

where \mathbf{v} is the gas velocity; the different metal-enriching sources from AGB winds ($\dot{X}_{i,*}$), SNIa's ($\dot{X}_{i,Ia}$), and SNI-I's ($\dot{X}_{i,II}$) are evidenced; star formation $\dot{X}_{i,SF} = (X_i/\rho) \cdot \dot{\rho}_{SF}$, where ρ is the gas density and $\dot{\rho}_{SF}$ is the gas sink due to star formation, is treated as a sink of local metals; and $\dot{\mathbf{m}}_{Q,i} = (X_i/\rho) \cdot \dot{\mathbf{m}}_Q$, where $\dot{\mathbf{m}}_Q$ is the momentum density for the gas that becomes Toomre-unstable (see Gan et al. 2019a for more details).

The simulations start at an age of 2 Gyr; therefore, the SSP contributes metals only via AGB winds and SNIa's ejecta. The weakly time-dependent metal input from AGB winds was calculated with “CELib,” an open-source software library for chemical evolution (Saitoh 2017). The abundance of the SSP is $Z_* = 1.5Z_\odot$ (where $Z_\odot = 0.0134$,

Asplund et al. 2009), as appropriate for a massive elliptical such as the one modeled (Thomas et al. 2010). The nucleosynthetic yields of SNIa are those of the N100 model of Seitenzahl et al. (2013), dominated by Fe and Si. The SNII yields are those of Nomoto et al. (2013), and average values are calculated for the progenitor's masses born with the adopted IMF of the new stars (see Gan et al. 2019a for more details).

The CGM infall provides low-metallicity gas with $Z_{CGM} = 0.002$ (i.e., $Z_{CGM} = 0.15 Z_\odot$). This gas accretes through the galaxy outskirts and produces a dilution of the abundance of the ISM of internal origin as it falls into the galaxy and mixes with the ISM. It does not appear explicitly in Equation 3 as the CGM infall is implemented as an outer boundary condition.

3 | RESULTS

We present the resulting chemical evolution of the ISM for an E2 elliptical galaxy of stellar mass at the present epoch of $M_* = 3.3 \times 10^{11} M_\odot$, effective radius $R_e = 7$ kpc, and central stellar velocity dispersion of 280 km s^{-1} . A central MBH and a spherical dark matter halo are also added, and the ordered rotational velocity of the stellar component is given by the Satoh decomposition, with exponentially declining rotation parameter k (Gan et al. 2019a). The presence of streaming velocity results, for the ISM, in the formation of a central cold disk, where star formation takes place; the latter then becomes an important metal-enriching source, mostly by SNe II. In fact, the ISM evolution undergoes recurrent cycles, during which the gas accumulates in the disk until the Toomre instability sets in, allowing for star formation and mass inflow from the disk toward the MBH. Thus, with a short delay, star formation is followed by accretion of disk material onto the MBH, whose feedback action via biconical broad absorption line (BAL) winds causes the ejection of recent metal-enriched gas from the disk (Figure 2). The abundance of the BAL winds is evaluated by tracking the chemical composition of the mass accreted via the inner boundary; the metals passing through the center are present in the BAL winds and injected back into the galaxy from its nucleus. Freshly produced metals are spread out at large distances (of $\simeq 1$ kpc and more); the abundance of the high velocity, outward-moving gas, can be many times the solar value. Note that these high abundances are fundamental for the gas surrounding the MBH to be multiphase and include a stable gas phase representing the broad line region (BLR) clouds (Chakravorty et al. 2009).

Figure 2 shows the density and temperature of the ISM during an outburst. The cold disk, extending for $r \simeq 150$




FIGURE 2 Meridional sections of the ISM density (left), temperature (middle), and abundance Z of all metals (right) for the ISM in the galaxy model described in Section 3 during a nuclear outburst. The cold disk is evident, as is the increase in Z above and below it, produced by the AGN wind. ISM, interstellar medium.




FIGURE 3 Map of the emission-weighted and -projected abundance of Fe (left) and Mg (right) for the ISM in the galaxy model described in Section 3; the snapshot is taken during a nuclear outburst, at the same time as Figure 2; note the linear scale instead of the logarithmic one used in Figure 2. ISM, interstellar medium.

pc, is clearly visible, as is the biconical wind structure in the temperature map. Note that the size of the cold disk strongly depends on the galaxy rotation profile. In the right panel, the total abundance Z is shown (the logarithmic scale on the axes emphasizes the central region). SNII products, transported by the winds, contaminate the region above and below the disk; they show many subsequent ejection episodes closely spaced in time, which take place during the nuclear outburst. As an example, in the outflowing region affected by the BAL wind with a radial velocity $v_r \geq 10^3$ km s⁻¹, the mass-weighted average abundance is $4Z_\odot$ (with a weighting toward products from SNII).

Figure 3 shows the emission-weighted and -projected maps for the abundance of Fe and Mg, which occurred at the same time as that shown in Figure 2 during the outburst. Interestingly, these two metal species show an opposite behavior. In fact, star formation increases the overall Z of the ISM, but in the SNII ejecta, the Fe is less abundant than in the mass lost by the old SSP. This causes Fe to be more abundant over the main galaxy body and less abundant in the region affected by recent star formation (the central disk and the biconical region affected by the spreading of metals due to the BAL winds). The symmetric behavior is shown by Mg, which is more abundant in the regions affected by SNII ejecta. Thus, SNII




FIGURE 4 Emission-weighted, -projected, and -circularized abundance of the ISM $Z(R)$, for the CGM₂₀ (in black) and CGM₇ (in red) models, and for the galaxy described in Section 3, at an age of 12 Gyr. Also shown (in blue) are the abundances Z_{old} at two epochs for the mass ejected by AGB winds + SNIa’s (with $Z_{\star} = 1.5Z_{\odot}$ for the SSP). The Z_{old} values are independent of the galaxy stellar mass, provided that the same IMF and Z_{\star} apply (see Section 3 for more details). ISM, interstellar medium; IMF, initial stellar mass function.

products such as Mg are a tracer of the distances reached by the ISM, driven by the AGN winds. Note that the biconical region in Figure 2 comes from the subsequent action of many outbursts during the galaxy’s lifetime and that each outburst produces an abundance pattern rich in substructure, which cannot be shown here due to the limited space.

Figure 4 shows the emission-weighted, -projected, and -circularized abundance profile, $Z(R)$, for the ISM when the SSP age is 12 Gyr. Also shown is the abundance of the mass contributed by the SSP sources (AGB + SNIa’s; hereafter “passive” abundance Z_{old}). As the SSP is assumed to be homogeneous across the galaxy, Z_{old} is position-independent; it depends instead on the SSP age as the metal injection of AGB winds and of SNIa follows different time dependencies (Section 2.1). In Figure 4, the values of Z_{old} at two ages of the SSP (2 and 13 Gyr) are shown. It can be noticed that (a) Z_{old} varies in a narrow range,¹ from 2.1 to 2.3 Z_{\odot} , over the whole time span, and (b) $Z(R)$ of the ISM when the SSP is 12 Gyr old (black line) is close to $Z_{\text{old}}(13 \text{ Gyr})$; $Z(R)$ is just slightly above it due to the enrichment contributed by new stars and does not show pronounced radial trends. The black line in the figure refers to the CGM₂₀ model, the same shown in Figures 2 and 3; the red line refers instead to its CGM₇ version, with a larger CGM infall (Section 2.1).




FIGURE 5 Emission-weighted, -projected, and -circularized Fe abundance of the ISM, $\text{Fe}(R)$, for the CGM₂₀ (in black) and CGM₇ (in red) models, and for the galaxy described in Section 3, at an age of 12 Gyr. Also shown (in blue) is Fe_{old} at two epochs, for the mass ejected by AGB winds + SNIa’s (with $\text{Fe}_{\star} = 1.5\text{Fe}_{\odot}$ for the SSP). Green squares show $\text{Fe}(R)$ for NGC507, a dominant elliptical galaxy in a small group, from XMM-Newton observations (Kim & Fabbiano 2004). See Section 3 for more details. ISM, interstellar medium.

The CGM₇ model shows a large radial gradient: Its $Z(R)$ is comparable to that of CGM₂₀ in the central galactic region but decreases outside of $R \sim 15$ kpc. The abundance of the ISM averaged *over the whole galaxy* is 2.4 Z_{\odot} for the CGM₂₀ model, dropping to $\sim 1.5Z_{\odot}$ for the CGM₇ one.

Figure 5 shows the emission-weighted, -projected, and -circularized Fe abundance profile, $\text{Fe}(R)$, for the ISM when the SSP age is 12 Gyr, again with the “passive” Fe abundance (Fe_{old}) at 2 and 13 Gyr of age for the SSP, and the CGM₇ model is included for comparison. The same trends of Figure 4 are apparent, with one difference: $\text{Fe}(R)$ stays below $\text{Fe}_{\text{old}}(13 \text{ Gyr})$, even at the center. This is explained by the fact that Fe_{old} is time-increasing, and the simple stellar population (SSP) age for the plotted ISM is 12 Gyr, and by the large difference in the Fe production between the SPP and the new stars (see the discussion above for Figure 3). Figure 5 also shows the observed $\text{Fe}(R)$ for a giant elliptical galaxy: Except for the innermost point, the outer region of $R \approx 20$ kpc, the agreement with the modeled ISM is good.

4 | CONCLUSIONS

In the MACER code for the investigation of the evolution of the ISM of elliptical galaxies, at high spatial resolution and with AGN feedback, the metal enrichment, transportation, and dilution processes have recently been implemented (Gan et al. 2019a). From these recent simulations for a massive elliptical, here, we have emphasized the following results:

¹The “passive” values in Figure 4 refer to the Kroupa IMF, but they would change very little for the Salpeter IMF.

1. The average Z of the hot ISM on the whole galactic scale is $\simeq 2.4Z_{\odot}$; this value can decrease to $\sim 1.5Z_{\odot}$ when the CGM infall rate is closer to the largest estimates from cosmological simulations.
2. For $R < 15$ kpc, $Z(R)$ of the hot ISM is close to the “passive” prediction (i.e., the abundance of AGB winds + SNIa’s), independent of the cosmologically motivated CGM infall rate. A small increase due to new star formation is present; this increase is disk-dependent and is thus related to the streaming velocity field of the host galaxy. The outer region of $R \simeq 15$ kpc, $Z(R)$, sharply decreases for the larger CGM infall rate.
3. The Fe abundance profile, $\text{Fe}(R)$, follows the same trends shown by $Z(R)$, with the difference of being *lowered* by the presence of new star formation. This feature is also shown in the abundance map of Fe and is contrary to that shown in the Mg abundance map.
4. During outbursts, the outflowing regions, driven by the BAL wind, can reach an abundance of $Z \simeq 4Z_{\odot}$ and can transport metals (mostly SNII products made in the disk) out to large radii. This finding is in agreement with observations that Z of a quasar BLR in massive galaxies is 0.3–1.0 dex larger than Z of the host galaxies and does not evolve with cosmic time, which implies that the metal enrichment is due to recent star formation, as in the simulations (Nagao et al. 2006; Xu et al. 2018).

While the above results are encouraging for the part concerning the role of AGN winds, the observed Z (and Fe) of the hot ISM demonstrate more problematic issues. Observed values are generally solar, or slightly supersolar, except for a few cases (one is shown in Figure 5), as established even recently, with the improved quality of both *Chandra* and *XMM-Newton* observations and data analysis (Humphrey & Buote 2006; Loewenstein & Davis 2010; Mernier et al. 2017, 2018). The observed Z , especially Fe, are lower than that predicted by models, which represents an old problem that has been alleviated in more recent investigations but still clearly persists. Contributing factors to it include residual uncertainties in the derived abundances, for example, due to the degeneracy of metallicity and emission measure and the multitemperature thermal structure of the ISM (de Plaa et al. 2017; Kim 2012); dust depletion of metals (Lakhchaura et al. 2019; Panagoulia et al. 2013); and incomplete mixing of SNIa’s ejecta (Matsushita et al. 2000; Tang & Wang 2010). Based on the present results, the CGM infall can also have an important effect. Another possibility is the mixing and stirring of metals between the inner and outer parts of the galaxies, as would be provided by a larger feedback wind efficiency, and/or infall of satellites. Finally,

we emphasize that the depletion of metals onto dust has not been inserted in the models yet, and thus, elemental abundances correspond to the total (dust plus gas phase) mass in that element. The depletion of refractory elements onto dust grains may be a large correction (Hensley et al. 2014), and this subject should be the focus of future work.

REFERENCES

- Arav, N., Borguet, B., Chamberlain, C., Edmonds, D., & Danforth, C. 2013, *MNRAS*, 436, 3286.
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *ARA&A*, 47, 481.
- Bartko, H., Martins, F., Trippe, S., et al. 2010, *ApJ*, 708, 834.
- Brennan, R., Choi, E., Somerville, R. S., Hirschmann, M., Naab, T., & Ostriker, J. P. 2018, *ApJ*, 860, 14.
- Carniani, S., Marconi, A., Maiolino, R., et al. 2015, *A&A*, 580, 102.
- Chakravorty, S., Kembhavi, A. K., Elvis, M., & Ferland, G. 2009, *MNRAS*, 393, 83.
- Ciotti, L., Pellegrini, S., Negri, A., & Ostriker, J. P. 2017, *ApJ*, 835, 15.
- Fabbiano, G. 2012, Hot interstellar matter in elliptical galaxies. in: *Astrophysics and Space Science Library*, eds. D.-W. Kim & S. Pellegrini, Vol. 378, Springer (Berlin), 1.
- Forbes, D. A., Alabi, A., Romanowsky, A. J., Kim, D.-W., Brodie, J. P., & Fabbiano, G. 2017, *MNRAS*, 464, 26.
- Gal, Z., Choi, E., Ostriker, J. P., Ciotti, L., & Pellegrini, S. 2019a, *ApJ*, 875, 109.
- Gal, Z., Ciotti, L., Ostriker, J. P., & Yuan, F. 2019b, *ApJ*, 872, 167.
- Goulding, A. D., Greene, J. E., Ma, C.-P., et al. 2016, *ApJ*, 826, 167.
- Hensley, B. S., Ostriker, J. P., & Ciotti, L. 2014, *ApJ*, 789, 78.
- Humphrey, P. J., & Buote, D. A. 2006, *ApJ*, 639, 136.
- Jiang, Y.-F., & Goodman, J. 2011, *ApJ*, 730, 45.
- Kim, D.-W. 2012, Hot interstellar matter in elliptical galaxies. in: *Astrophysics and Space Science Library*, eds. D.-W. Kim & S. Pellegrini, Vol. 378, Springer (Berlin), 121.
- Kim, D.-W., & Fabbiano, G. 2004, *ApJ*, 613, 933.
- Lakhchaura, K., Mernier, F., & Werner, N. 2019, *A&A*, 623, 17.
- Loewenstein, M., & Davis, D. S. 2010, *ApJ*, 716, 384.
- Lu, J. R., Do, T., Ghez, A. M., Morris, M. R., Yelda, S., & Matthews, K. 2013, *ApJ*, 764, 155.
- Matsushita, K., Ohashi, T., & Makishima, K. 2000, *PASJ*, 52, 685.
- Mernier, F., de Plaa, J., Kaastra, J. S., et al. 2017, *A&A*, 603, A80.
- Mernier, F., Werner, N., de Plaa, J., et al. 2018, *MNRAS*, 480, L95.
- Nagao, T., Marconi, A., & Maiolino, R. 2006, *A&A*, 447, 157.
- Negri, A., Posacki, S., Pellegrini, S., & Ciotti, L. 2014, *MNRAS*, 445, 135.
- Nomoto, K., Kobayashi, C., & Tominaga, N. 2013, *ARA&A*, 51, 457.
- Panagoulia, E. K., Fabian, A. C., & Sanders, J. S. 2013, *MNRAS*, 433, 3290.
- Pellegrini, S. 2012, Hot interstellar matter in elliptical galaxies. in: *Astrophysics and Space Science Library*, eds. D.-W. Kim & S. Pellegrini, Vol. 378, Springer (Berlin), 21.
- Pellegrini, S., Ciotti, L., Negri, A., & Ostriker, J. P. 2018, *ApJ*, 856, 115.
- de Plaa, J., Kaastra, J. S., Werner, N., et al. 2017, *A&A*, 607, 98.
- Saitoh, T. R. 2017, *AJ*, 153, 85.
- Seitzahl, I. R., Ciaraldi-Schoolmann, F., Roepke, F. K., et al. 2013, *MNRAS*, 429, 1156.
- Tang, S., & Wang, Q. D. 2010, *MNRAS*, 408, 1011.

- Thomas, D., Maraston, C., Schawinski, K., Sarzi, M., & Silk, J. 2010, *MNRAS*, 404, 1775.
- Tombesi, F., Melendez, M., Veilleux, S., Reeves, J. N., Gonzalez-Alfonso, E., & Reynolds, C. S. 2015, *Nature*, 519, 436.
- Werner, N., McNamara, B. R., Churazov, E., & Scannapieco, E. 2019, *SSRv*, 215, 5.
- Xu, F., Bian, F., Shen, Y., Zuo, W., Fan, X., & Zhu, Z. 2018, *MNRAS*, 480, 345.

AUTHOR BIOGRAPHY

Silvia Pellegrini is an associate professor at the Department of Physics and Astronomy of the University of Bologna, where she earned a PhD in Astronomy in 1993. She is an Astrophysicist with main interests

in X-ray astronomy. Her work has focussed on normal and active galaxies, from an observational point of view (using data obtained with the satellites Chandra and XMM-Newton), and a theoretical one (using numerical simulations of hydrodynamics, and dynamical models of galaxies).

How to cite this article: Pellegrini S, Gan Z, Ostriker JP, Ciotti L. Metal abundances in the MACER simulations of the hot interstellar medium. *Astron. Nachr.* 2020;341:184–190. <https://doi.org/10.1002/asna.202023776>